

Optimal Filtering

Exercise 1

Gustaf Hendeby*
gustaf.hendeby@liu.se

Version: 2023-11-11

Problem 1.1 Consider the following block matrix

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix},$$

where $|A| \neq 0$, $|D| \neq 0$, and $|A|$ denotes the determinant of the matrix A .

a) Derive the block UDL factorization of

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix},$$

i.e., find X , Y , Δ_1 , and Δ_2 such that

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} I & X \\ 0 & I \end{pmatrix} \begin{pmatrix} \Delta_1 & 0 \\ 0 & \Delta_2 \end{pmatrix} \begin{pmatrix} I & 0 \\ Y & I \end{pmatrix}.$$

b) Derive the inverse of

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix}.$$

c) Show that

$$\left| \begin{pmatrix} A & B \\ C & D \end{pmatrix} \right| = |A - BD^{-1}C| |D|.$$

Problem 1.2 Let

$$\begin{pmatrix} x \\ y \end{pmatrix} \sim \mathcal{N} \left(\begin{pmatrix} \mu_x \\ \mu_y \end{pmatrix}, \begin{pmatrix} \Sigma_{xx} & \Sigma_{xy} \\ \Sigma_{yx} & \Sigma_{yy} \end{pmatrix} \right).$$

Show that the conditional pdf $p_{X|Y}(x|y)$ is Gaussian with mean

$$\mu_{x|y} = \mu_x + \Sigma_{xy} \Sigma_{yy}^{-1} (y - \mu_y)$$

and covariance

$$\Sigma_{x|y} = \Sigma_{xx} - \Sigma_{xy} \Sigma_{yy}^{-1} \Sigma_{yx}.$$

Hint: Use the results from Problem 1.1.

*These exercises are heavily inspired by exercises used by Isaac Skog (LiU) and Mats Bengtsson (KTH).

Problem 1.3 Suppose X and Y are jointly distributed random variables. When Y is unknown, an appealing estimate of the value of X is its mean value $\mu_x = E[X]$. This estimate has the property

$$E[\|X - \mu_x\|^2] \leq E[\|X - z\|^2]$$

for all z , and has the average error $E[\|X - \mu_x\|^2]$. Now suppose that one is told that $Y = y$. Let $\hat{x} = E_{X|Y=y}[X]$. Show that

$$E_{X|Y=y}[\|X - \hat{x}\|^2] \leq E_{X|Y=y}[\|X - \mu_x\|^2].$$

This directly gives the intuitively reasonable result that the mean estimation error $E[\|X - \hat{x}\|^2]$ averaged over all values of X and Y is bounded from above by $E[\|X - \mu_x\|^2]$. When is this bound attained, *i.e.*, when is there no improvement in the knowledge of X given the value of Y ?

Problem 1.4 Given a zero-mean wide-sense stationary process $y(t)$ with *auto correlation function* (acf) $r_{yy}(\tau)$, derive the linear least-squares estimate of

$$\int_0^T y(t) dt$$

in terms of its values at the end points $y(0)$ and $y(T)$.

Assuming that the covariance function is

$$r_{yy}(\tau) = e^{-\alpha|\tau|},$$

show that the estimate is

$$\frac{1}{\alpha} \tanh \frac{\alpha T}{2} (y(0) + y(T)).$$

Furthermore, provide an approximate estimate that holds for small T .

Problem 1.5 Assume X and Z are independent (jointly) Gaussian random variables with means μ_x and μ_z , respectively, and with covariance matrices Σ_{xx} and Σ_{zz} . Then $Y = X + Z$ is Gaussian and $p_{X|Y}(x|y)$ is a Gaussian density function. Show that the associated conditional mean and covariance are

$$\begin{aligned} \mu_{x|y} &= \Sigma_{zz}(\Sigma_{xx} + \Sigma_{zz})^{-1}\mu_x + \Sigma_{xx}(\Sigma_{xx} + \Sigma_{zz})^{-1}(y - \mu_z) \\ &= \Sigma_{x|y}(\Sigma_{xx}^{-1}\mu_x + \Sigma_{zz}^{-1}(y - \mu_z)) \end{aligned}$$

and

$$\begin{aligned} \Sigma_{x|y} &= \Sigma_{xx} - \Sigma_{xx}(\Sigma_{xx} + \Sigma_{zz})^{-1}\Sigma_{xx} \\ &= (\Sigma_{xx}^{-1} + \Sigma_{zz}^{-1})^{-1}, \end{aligned}$$

respectively.

Hint: Begin by deriving the joint density $p_{X,Y}(x,y)$ and assume that the inverses exist.